1 □□□□□□□□□□

□□□□□

VITAFOL H Hardener

□□□□□□□□□

□□□□□□□:

□□□□□□□

H226	□□□□□□□□□□□
H319	□□□□□□
H332	□□□□□□□□
H335	□□□□□□□□□□□□□
H361d	□□□□□□□□□□□□
H372	□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□

P210	□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□-□□□
P260	□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□
P280	□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□
P337+P313	□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□□□

□□□□□□□□□

3 □□□□□□□□□

□□□

□□□□□□

CAS□□	□□□	□□
	□□ (EC □□ No. 1272/2008)	
78-10-4	ethyl silicate, tetraethyl silicate	80 - < 85 %
	Flam. Liq. 3, Acute Tox. 4, Eye Irrit. 2, STOT SE 3; H226 H332 H319 H335	
93925-43-0	Silicic acid (H4SiO4), tetraethyl ester, reaction products with bis(acetyloxy)dioctylstannane	10 - < 15 %
	Flam. Liq. 3, Repr. 2, Acute Tox. 4, Eye Irrit. 2, STOT RE 1, Aquatic Chronic 4; H226 H361d H302 H319 H372 H413	
68299-15-0	Bis(neodecanoyloxy)dioctylstannane	1 - < 5 %
	STOT RE 2, Aquatic Chronic 4; H373 H413	

H□□□EUH□□□□□: 16□□□□□□□□□ □

SCL□M-factor □□/□□ ATE

CAS□□	EC□□	□□□	□□
	SCL□M-factor □□/□□ ATE		
78-10-4	201-083-8	ethyl silicate, tetraethyl ester	80 - < 85 %
	□□: □□□□□□□ = 11 mg/l (□□); □□: □□□□□□□ = 1.5 mg/l (□/□□□); □□: LD50(50%□□□) = 5860 mg/kg		
93925-43-0	300-346-5	Silicic acid (H4SiO4), tetraethyl ester, reaction products with bis(acetyloxy)dioctylstannane	10 - < 15 %
	□□: □□□□□□□ = 500 mg/kg		

4 □□□□

□□□□□□□□□□□

□□□□□□□

□□□□□□□□□□□□□ □□□□□□□□□□□□□

□□□□□□□□□

□□□□□□□□□

□□□□□□□□□

□□: □□□

□□□□: □□□□□□□□□

□□□□: □□□□□□□□□

□□□□□□

□□: □□□

□□: □□□

□□□□: □□□□□□□□□

□H□: □□□□□□□□□

□□□: □□□

□□□□□□□□□□

□□□□□□□□□

n-□□□□□□□/□□□□□□: □□□□□□□□□

□□□: <=1100 hPa
(□ 50 °C)

□□: □□□□□□□□□

□□□□□□□: □□□□□□□□□

□□□□□□□

□□□□□□□□□□□□□□□□□□□□

□□□□

□□□□□□□□□ □

□□□□□□□□□□

□□□□□: 0,0 %

□□□□: □□□□□□□□□

□□□□□

10 □□□□□□□□□□

□□□

□□□.

□□□□□□□

□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□□□□□□□□

□□□□□□□□□

□□□□□□□□□□□□□□□□□□□

CAS□□	□□□				
	□□□□□	□□□	□	□□□□□	□□
78-10-4	ethyl silicate, tetraethyl silicate				
	□□□	LD50(50%□□□) 5860 mg/kg			
	□□□□□ □□	□□□□□□□ 11 mg/l			
	□□□□□ □/□□□	□□□□□□□ 1.5 mg/l			
93925-43-0	Silicic acid (H4SiO4), tetraethyl ester, reaction products with bis(acetyloxy)dioctylstannane				
	□□□	□□□□□□□ 500 mg/kg			

□□□□□□□□□

□□□□□□

□□□□□/□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□□□□□□/□□□□/□□□□□□

□□□□□□□□□□□ (Silicic acid (H4SiO4), tetraethyl ester, reaction products with bis(acetyloxy)dioctylstannane)

□□□□□□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□□□□□□□□□□□

□□□□□□□□□□□□□□ (ethyl silicate, tetraethyl silicate)

□□□□□□□□□□□□□□□□

□□□□□□□□□□□□□□□□□□□□□□□□□□ (Silicic acid (H4SiO4), tetraethyl ester, reaction products with bis(acetyloxy)dioctylstannane)

□□□□□□□□□□

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□EC□□No 1272/2008 [CLP]□□□□□□□□□□□□□□□□□□□□□□

12 □□□□□□□

□□

□□□□□□□□□□□□: □□□□□□

□□□□□□□□

□□□□□□□□□□□□□□□□□

□□□□□□

□□□□□□□□□□□□□□□□□

□□□□□□□□

□□□□□□□□□□□□□□□□□

□□□□□□□□□

□□□□□□□□□□□□□□□□□□□

13 □□□□□□□

□□□□□□□□

□□□□□

□□□□□□□□□□□□□□□□□□□□□□□□□ □□/□□□□□□□□□□□□□□□ □□□□□□□□□□□□□□□□□□□□□

□□□□□□□

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□. □□□□□□□□□□□□□□□□□□□□□□□□□□□

14 □□□□□□□

□□□□ **(IMDG)**

UN□□□□□□ID□□:	UN 1292
□□□□□□□□□□:	TETRAETHYL SILICATE
□□□□□□□□□□□□□□□□:	3
□□□□□□□□□:	III
□□□□□□□:	3

□□□□□□:	-
□□□□:	5 L
□□:	E1
EmS:	F-E, S-D

□□ **(ICAO-TI/IATA-DGR)**

UN□□□□□□ID□□:	UN 1292
□□□□□□□□□□:	TETRAETHYL SILICATE
□□□□□□□□□□□□□□□□:	3
□□□□□□□□□□:	III
□□□□□□□□:	3

□□□-□□:	10 L	
Passenger LQ:	Y344	
□□:	E1	
IATA□□□□□□-□□:		355
IATA□□□□□-□□:		60 L
IATA□□□□□□□□□□:		366
IATA□□□□□□□□□□:		220 L

□□□□□□□□□

□□□□□□□□□: □□□

□□□□□□□□□□□□□□□□

□□: □□□□□□□

MARPOL 73/78 □□□□□□ II □□□□ IBC Code □□□□□□□□□□□□

□□□

15 □□□□□

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□/□□

□□□□□□

□□□□: □□□□□□□□□□□□□□□□□
□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□□□□□□□□□□□: 2 - □□□□□□□

16 □□□□□□□

□□

□□□□□□□□□□□□□□□□□□□